


Climate Change in the Oregon 5th Senate District

August 2019

Updated May 2023



History, Projections, and Consequences

1. The temperature has risen some 1°F during the last half of the 20th Century and may rise 8°F by the end of the 21st Century.
2. While precipitation has been steady and is likely to remain so annually, wetter winters and drier summers.
3. The trend of declining snowfall will continue through the century, possibly dropping to 10% of historic levels by 2100.
4. These precipitation projections, combined with the trend towards increasing heavy rainfall and reducing light rainfall will likely increase flooding and compromise irrigation availability in those months when it is most needed.
5. Several important forest species both commercially and in terms of forest composition will likely be compromised as climate change overtakes the District.
6. The western wildfire season is already 105 days longer than in the 1970s, while reduced snowpack, warmer summers and earlier snowmelt will increase wildfire risk, with 200 – 300% of the area burned by mid-century.
7. Agricultural activities such as wine growing that depend on temperature and water are likely to be threatened through the century.
8. As sea levels rise, increased urban storm damage and destruction will be probable in addition to the loss of beaches and coastal wetlands.
9. Those engaged in agriculture, forestry or fisheries will be most affected by the forthcoming climate trends, will need most to adapt, and probably should be most supportive of mitigation efforts.
10. The main climate impacts to health are likely: storms, floods, and sea level rise. The main health concerns will be: disruption in core services, injuries, displacement, landslides, income loss, economic instability, and mental health impacts. Vulnerable communities will be: low-income households, older adults, people living on steep slopes, farmers of fish and shellfish, first responders, and children and pregnant women.
11. To achieve required emissions reduction goals, we need to reduce emissions 45% below 2010 levels by 2030; this requires immediate action!

Compiled by Emily Patrick & Alan Journet (patricke@sou.edu, 541-880-6563)
(alanjournet@gmail.com, 541-301-4107)

Updated by Hogan Sherrow (hogan@you-evolving.com, 541-415-1013)

For a more complete summary, including sources, from which these points are taken, visit: <http://socan.eco/oregon-legislative-districts/>

We invite copying of these materials, but request that authorship together with the SOCAN logo and attribution be retained.

Please contact Alan Journet (alan@socan.eco 541-301-4107 or 541-500-2331) with any queries.

Climate Change in the Oregon 5th Senate District

Compiled by Emily Patrick & Alan Journet

(patricke@sou.edu, 541-880-6563)

(alanjournet@gmail.com, 541-301-4107)

August, 2019

Updated by

Hogan Sherrow

(hogan@you-evolving.com, 541-415-1013)

May 2023

Global and Regional Temperature:




Figure 1. Historic global temperature trend (NASA 2023).




Figure 2. Historic U.S temperature trend. (NOAA 2023).

Data from NASA and NOAA reveal that the Global and U.S. atmospheric temperatures have increased substantially since 1880 (Figures 1 and 2) with the greatest effect occurring in the last five decades.




Figure 3. Global temperature projections to 2100 relative to the 1850-1900 average. (IPCC 2021)

the human behavior that leads to specific atmospheric greenhouse gas concentrations. The SSP5-8.5 pathway incorporates (SOS 2022) a: “push for economic and social development ... coupled with the exploitation of abundant fossil fuel resources and the adoption of resource and energy intensive lifestyles around the world.” Effectively this seems to echo the RCP8.5 projections employed in the previous IPCC report and is the trajectory we are currently following globally. This scenario would likely result in global temperatures in the range of 3 to 5.1°C (5.4 to 9.18°F) above pre-industrial revolution temperatures by 2100 (Figure 3).

Meanwhile, projections further into the future have been provided by the Intergovernmental Panel on Climate Change (IPCC) in terms of RCP scenarios (Figure 4) The RCP 2.6 scenario

assumes we rapidly eliminate emissions, whereas RCP 8.5 assumes we follow the current trajectory of accelerating emissions. RCP 6.0 and 4.5 assume intermediate trajectories of emissions between the extremes. Note that only the RCP2.6 scenario results in a long-term global temperature increase below 2°C above pre-industrial conditions - the upper target for the 2015 Paris Agreement. Because the actual temperature trajectory we have experienced follows the RCP 8.5 scenario this has been dubbed the Business-As-Usual (BAU) scenario; we have yet to undertake sufficient actions globally to slow this trend.




Figure 4. Long term global temperature trends according to RCP values. (Jones 2017).

Meanwhile, temperature projections for this century in Oregon (Fleishman 2023, Figure 5) suggest a similar range of temperature increases possibly reaching over 13°F above the 1970-1999 average by the end of the century under the BAU scenario (RCP 8.5).




Figure 5. Oregon temperature history and projections through the century; baseline: 1970 – 1999 (Fleishman 2023)

Whether we consider the global or Oregon future, the higher range of temperature increase would be unmanageable. It would devastate natural systems (see below:) and simultaneously threaten our climate dependent agricultural, ranching, and forestry activities. Bark beetle and other pest destruction of forests would likely increase as warmer temperatures enhance insect growth and development rates and enable larger

overwintering populations. Similarly, invasion of natural and agricultural systems by drought tolerant invasive species and pests will likely be enhanced.

Regional Precipitation:




Figure 6. History and projections for precipitation statewide. Fleishman 2023

Annual precipitation is expected to increase very slightly (if at all) in Oregon through the balance of this century (Figure 6). However, the 2018 US Climate Change Assessment Report (Easterling *et al.* 2017) provides projections for seasonal late century precipitation patterns (Figure 7) according to the ‘business as usual’ RCP 8.5 scenario.

The region generally is expected to exhibit fall and spring seasons that are little different from historical patterns, with winters possibly a little wetter. Notably, however, accentuating the Mediterranean 'winter wet - summer dry' climate, winters will be wetter, and summers will likely be drier.

Evaporation caused by increasing temperature will likely counter any increase in precipitation such that drought conditions continue. Water resources, already severely compromised in many locations, will become more threatened as snowpack continues to decline. Meanwhile, the current trend of precipitation occurring more frequently as severe storms rather than the light drizzle that replenishes soil moisture will continue. This will likely increase the risk of floods, soil erosion and landslides.




Figure 7. Projected change in precipitation to 2077-2090 compared to 1960-2005 average; stippled areas indicate large change compared to natural variation; hatched areas small change. (Easterling *et al.* 2017)




Figure 8. Potential increase in fire risk from the 1971-2000 baseline across the U.S. assuming the RCP 8.5 scenario. Gao *et al.* 2021

Stream and river flow occurring during summer/fall will decline and become warmer compromising many iconic Pacific Northwest cold-water aquatic species. Meanwhile, peak river flow will continue to advance earlier in the year, even into late fall.

Gao *et al.* (2021) depicted the increasing risk of fire across the nation (Figure 8) under the RCP 8.5 scenario indicating that this would likely lead to increased fire probability throughout most of Oregon of at least 50%.

Several years ago, the national climate assessment, (Melillo *et al.* 2014) reported the impact of increasing temperature of just 2.2°F on area burned from wildfire, a condition potentially arriving by mid-century (Figure 9). The range in increase is from 100% meaning a doubling of the area burned to 700% meaning 8-times the current area.

Mote *et al.* (2019) presented a summary of the potential increase in extreme fire risk days by mid-century (2040 - 2069) compared to historical conditions (1971 - 2000) where an increase of up to 14 days in the SE corner of the state is evident (Figure 10).

The fire season, already extended by 105 days since 1970s (Kenward *et al.*




Figure 9. Anticipated wildfire consequences of a 2.2°F warming in area burned (Melillo *et al.* 2014).




Figure 10. Increase in number of days experiencing high fire risk by mid- century (2040 - 2069) compared to the end of the last century (1971-2000). Mote *et al.* 2019.

2016), will likely become longer and more severe in Oregon. Even though our natural ecosystems have evolved with fire and are thus fire prone, fire adapted, and fire dependent, future trends may pose a serious threat to ecosystem ongoing health. In addition, of course, both human safety and human health will likely be threatened. It was recognized long ago (Westerling *et al.* 2006) that warming and early spring snowmelt correlate with increasing fire risk.

Natural System Consequences

That the geographic distribution of our natural ecological systems (biomes) is largely determined by the variables of temperature and precipitation (water availability) has been understood for decades. Community ecologist Robert Whittaker (1975) developed a classic chart indicating this relationship (modified in Figure 11). The chart depicts the climatic conditions that allow each of the designated biomes to exist. The critical message is that even a small shift in either of these variables from current conditions may threaten the viability of the biomes and the species of which they are comprised. This will be especially true for biomes currently existing at the edge of the climatic range that they require. It is especially worth




Figure 11. Global distribution of natural ecological systems (biomes) in relation to mean annual temperature and precipitation patterns. (Modified from Whittaker 1975.)

comparing these temperature ranges to the potential shifts in Oregon's temperature through the century (Figure 5) from which it is evident that most of our state's precious natural systems will be threatened, and some (especially high-altitude cool climate systems) will likely be eliminated under future conditions. It is worth noting, also, that the same variables control our agricultural productivity, and clearly, our forest viability. Thus, climatic shifts of the dimensions anticipated, absent any adjustment in our collective behavior that thwarts

that trajectory, will likely compromise agriculture and forestry throughout the state. Indeed, Dalton *et al.* (2017) indicate not only that "different trees have varying degrees of sensitivity to climate change and adaptive capacity." but also that "suitable climates for many important tree species and vegetation types may change considerably by the end of the 21st century...." Climate envelope projections (Rehfeldt and Crookston 2023), which assess the optimal conditions for tree species on the basis of their current and recent historic range and map these condition into the future, suggest that under the RCP 8.5 scenario, several species will likely suffer range reduction: Douglas fir, Western hemlock, Ponderosa pine, Grand fir, Western larch, Sugar pine, White fir, Pacific madrone, Western juniper, Western redcedar, Tanoak, and California laurel. Meanwhile, by the end of the century, the following species will likely find the Oregon climate completely outside their range (i.e., they will be extirpated from the state): Sitka spruce, Engelmann spruce, Lodgepole pine, Subalpine fir, and Jeffrey pine. Oregonians dependent on commercial timber harvest should be the first to demand climate action in the state.

Coastal Concerns:

Though much of Oregon is land-locked, and will suffer little directly because of ocean consequences, coastal regions and economies will have to contend with warming oceans, sea level rise, and increasing ocean acidification.

Warming Oceans. Although there is considerable seasonal fluctuation in ocean temperature, warming of oceans in the Northwest are already documented with a reported and anticipated increase at the rate of 0.35°C per decade (Alexander *et al.* 2018) off the coast of Oregon over

the period 1976 - 2009. Besides influencing species directly, temperature changes impact such events as algal blooms and shellfish poisoning.

Sea Level Rise. Sea levels are rising and will continue to rise for two reasons: First, water expands as it warms from 4°C (approximately 37°F). Thus, as the ocean warms, it inevitably expands, and sea level inevitably rises. Second, as land borne ice enters the ocean, whether as water or ice, it increases the volume of the ocean. Both these phenomena have already caused sea level to rise and are expected to continue this impact. The impact is influenced by the pattern of land adjustment: if land is rising, the impact is reduced, whereas subsiding coastal land will exacerbate the impact. This complication is particularly relevant to the impact of the Cascadia Subduction Zone (CSZ) where a rising or falling land tectonic plate will influence apparent and locally detected sea level rise along the coast. The impact of the oceanic Juan de Fuca plate sliding under the continental North American plate is a rising continental plate (Lieberman 2012) apparently confounding the ability of a land-based gauge to detect sea level rise. However, should the earthquake occur, there will likely result a drop in the land level of a meter (3 feet) or so. Mote *et al* (2019), however, indicate that by century's end, the actual sea level rise off the coast of Oregon could plausibly reach 8 feet, a value reiterated in Fleishman (2023). During storm surges, a higher sea level will generate conditions that promote far greater storm damage and flooding than would otherwise have been the case. The impact of Hurricane Sandy in 2012 was a perfect illustration of this problem. Not long ago, the suggestion that New York subways could be flooded by a coastal storm would not have been taken seriously – yet it happened! Results of ocean rise, such as increased erosion and compromised coastal habitat integrity for tidal flat, estuary, and marsh natural communities, could become serious.

Ocean Chemistry. Serious as direct climatic consequences are, they do not constitute the sum total of the impacts of our emitting carbon dioxide into the atmosphere.

Because carbon dioxide is absorbed by our oceans, and is transformed into carbonic acid, oceans are becoming more acidic. This is detrimental for marine organisms with carbon-based shells since either, they are unable to form shells in acidic conditions, or they lose shells already established. Bednaršek *et al.* (2020) demonstrated that ocean acidification off the coast of Oregon is already having a negative effect on Dungeness crab (*Metacarcinus magister*) shell formation and durability. Dungeness crab is one of the most valuable species on the Oregon coast, and the further acidification of our coastal waters could be catastrophic for this population. Additionally, acidosis, a build-up of acidic conditions in the tissues, threatens many marine life forms.

In addition, warming oceans exhibit reduced oxygen levels, potentially critical for marine animals since, like terrestrial animals, they rely on oxygen for basic metabolic respiration.

These consequences of increasing atmospheric greenhouse gases (notably carbon dioxide) pose threats to marine life, and thus to our fisheries, coastal economies, recreation, and tourism.

Rural vs Urban Oregon:

Rural communities are typically characterized by local economies and livelihoods that are reliant on direct interactions with the environment through agriculture, timber, fishing or outdoor based tourism activities. Urban communities, by contrast are typically characterized by local economies and livelihoods that are reliant on activities that do not include direct interactions with the environment. The result is that climate change has a far greater direct effect on rural communities than urban areas, including the direct effects of reduced snowpack, decreased river levels, rising seas, altered growing seasons, extended drought, increasingly hot summers, and increased wildfire risk. This has led to the misconception that urban communities are not vulnerable to the impacts of climate change.

While rural communities are on the frontlines of the climate crisis and some of the most vulnerable communities across Oregon, urban areas are also vulnerable. The heat related deaths in the Portland-metro area in the summer of 2020 and the Labor Day fires later that year demonstrated that urban areas are under direct threat from the impacts of climate change. Beyond the direct impacts of climate change, urban areas rely on healthy rural regions for their water supply, their agricultural, and forestry products and recreational activities in wild and less developed areas. Indirect effects, therefore, can be substantial. Climatic events that compromise natural systems, and thus urban watersheds, and regional agriculture and forestry will also have a profound impact on life in the urban centers. This impact will not be limited to impacts on prices of food and wood products but will also compromise regional recreational opportunities. Additionally, because the Pacific Northwest is projected to suffer less than most of the country from the warming climate, the region will become the target for climate refugees from across the U.S. These migrants will increase our population and place a greater burden on our natural resources and dwindling water supplies.

The 5th Oregon Senate District Climate History and Projections:

Although climate change is a complex issue, current models indicate several important trends in weather and climate that Oregon's 5th senate district is likely to experience if carbon emissions continue to increase. The trend for Lincoln County indicates an increase in mean annual temperature (Figure 12). Red represents the business as usual scenario of accelerating fossil fuel use and greenhouse gas emissions while blue represents a lower but

still increasing emissions trajectory. The temperature rose about 1°F during the latter half of the 20th century with a projected rise of up to a further 8°F above the mean for that period by the end of the century. Trends and projections for the other coastal counties of Senate District 5 are similar.




Figure 11. The historical and projected annual temperatures for Lincoln County (USGS 2021).

Precipitation history and projections from Lincoln County, presented in Figure 13, show essentially a constant history and future but with greater variability, meaning wetter and drier years. Should temperatures follow the path suggested above, however, this will likely lead to increased drought. As above, trends and projections for other coastal Senate District 5 counties are similar.




Figure 13. *Precipitation history and projections for Lincoln County, Oregon (USGS 2021).*

Meanwhile, snowfall data for the same county (Figure 14) reveal a recent history of decline and a projected future of continuing reduction. Since accumulated snowfall serves as a source for summer irrigation, the reduced snowfall will likely compromise agriculture in the coming century. Again, trends and projections for other coastal Senate District 5 counties are similar. The reduced snowpack also suggests more severe wildfire seasons are likely. The projected trend in snowpack is for a continued decline, possibly to only 10% of historical levels by late century.




Figure 14. Snowfall history and projections for Lincoln County, Oregon (USGS 2021).

Combined with the trend towards precipitation falling in heavy downpours on more days rather than light rain on many days as seen historically, this will likely result in earlier and decreased stream flow, a consequence that poses a serious threat to those agricultural activities dependent on late summer and early fall snowmelt as an irrigation source. The current trend towards precipitation falling in more frequent heavy thunderstorms as opposed to the light rainfall that rejuvenate soil moisture is also expected to continue. This means an increased risk of floods, soil erosion and landslides.

Federal Congressional District 4




Figure 15. Temperature trend for Federal Congressional District 4 (CCT 2021).

Oregon State Senate District 5 falls within Federal Congressional District 4. The historic temperature trend for this Congressional District (Figure 15) shows an increase of app 2.1°F per century since 1895 (a value comparable with this State Senate District) but note that the rate of increase itself has increased to 3.8°F since 1960.

Oregon 5th Senate District Economy:

Oregon's 5th senate district has a vibrant, viable and diverse economy that is based around agriculture, forestry, fishing, tourism, shipping, and wine growing. Unfortunately, all these industries will undoubtedly be negatively impacted by climate change in the next century if nothing is done now to mitigate potential problems before they arise.

Oregon's 5th senate district enjoys some of the region's best commercial timber; logging and timber production are important parts of the district's economy. However, predictions for the future of this industry under climate change scenarios are grim. As temperatures continue to increase over the next century, the range of many of Oregon's commercially important tree species is likely to shift. Projections for the future success of Douglas fir, Western hemlock and Sitka spruce, the most important local species, are presented in Figures 16 – 18. The 'Business as Usual' pathway is likely to compromise the viability of these species posing serious problems for the local timber industry. This would mean not only a loss of jobs for the region, but a loss of an important part of Oregon's timber producing heritage. In addition, given the ability of many Oregon forests to store carbon (Hudiburg *et al.* 2009), it is critical that climatic conditions not diverge such that these important species are compromised.

Figure 16. Douglas fir, *Psuedotsuga menzeisii* appropriate climate through the century (Crookston and Radtke 2023).




Figure 17. Western hemlock, (*Tsuga heterophylla*) appropriate climate through the century (Crookston and Radtke 2023).





Figure 18. Sitka spruce, *Picea sitchensis* appropriate climate through the century (Crookston and Radtke 2023).



In these graphs, purple represents climatic conditions ideal for the species, green and yellow represent sub-optimal conditions, while grey represents conditions unfavorable for the species. Those in the timber industry should understand that absent serious program to control greenhouse gas emissions, there will be no timber industry in Oregon later this century.

Furthermore, given the ability of many Oregon forests to store carbon (Hudiburg *et al.* 2009; Law *et al.* 2018), it is critical that climatic conditions not diverge such that these important species are compromised. Halofsky *et al.* (2016) discuss the potential and disturbing impacts of climate change of SW Oregon's forests.

AVERAGE GROWING SEASON TEMPERATURES THE RANGE IN THE ABILITY TO RIPEN VARIETIES Northern Hemisphere (Apr-Oct), Southern Hemisphere (Oct-Apr)




Figure 19. Grape varietal optimum growing season temperature (Jones 2015).

Potential Agricultural Impacts:

Our field crops are planted in soil and climatic conditions to which they are well adapted. This means adjustments from current climate can be detrimental. The agricultural 'one-degree problem' occurs because increasing temperature generally reduces crop yield. For each degree C temperature rise crop yield drops some 5 - 10% (Brown 2006). Meanwhile, the 'business as usual' scenario of increasing greenhouse gas emissions suggests that throughout Oregon the temperature will likely

increase 5 or more degrees C with decreasing soil moisture (USGS 2014) posing a great risk of extended drought. Farmers and home gardeners in Oregon should be concerned about a compromised future.

Future climate patterns as projected would negatively impact the economy through a reduction in crop yields since increasing temperature consistently reduces crop productivity and a potential for lost tourism due to wildfire. The blossoming wine industry and the pears produced by and for Harry and David would also be affected by the altered growing season. A potential problem for pear growers is the need for a solid winter chill period. This is decreasing. While not immediately a problem, if the trend of decreasing chill hours continues the consequences for pear production could become relevant. Legally grown marijuana and hemp have also become important components of the economy in Senate District 5. Both are water intensive practices and will, therefore be severely impacted by reduced precipitation and stream flows.

The predominant wine varieties in this area are Pinot Gris, Syrah, Merlot, Cabernet Sauvignon, Pinot Noir, and Chardonnay. Figure 18 depicts the growing season optimal temperatures for varieties grown in the region including the impact climate change will likely have on wine growing. While many of the grape varieties grown in this area seem reasonably well-adapted to mid-century growing season temperatures, even some of the warm climate varieties could be compromised by late century. However, of particular note are the cooler growing season varieties of the region (especially Illinois Valley wines) such as Pinot gris, and Gewürtstraminer, which could be severely compromised even by mid-century.

If climate trends continue as projected, Oregon's 5th Senate District will experience considerable natural and economic disruption. In order to sustain a vibrant economy, the region will find it necessary to adapt. Avoiding the worst-case scenario depicted in these projections will require the concerted effort of elected leaders at all levels of government, regional, national, and international.

Potential Health Risks:

Not only will climate change be negative for our economy, it will also change the lives of people in the 5th Senate district. This District and surrounding areas have become increasingly popular as retirement locations; Climate change and its consequences target the most vulnerable - such as the young and the old. The consequences depicted here could have a severe impact on the health of the elderly. Many of the health consequences involve respiratory problems for this vulnerable segment of the population. Heat waves and particulates emitted by wildfires can be particularly hazardous to those with respiratory problems.

According to the Oregon Health Authority (2020), the impacts of climate change on health are likely to be numerous: poor air quality, poor water quality, respiratory illness, occupational and recreational hazards, heat-related illness, residential displacement, contaminated drinking water, water insecurity, food insecurity, vector-borne disease, income loss, economic instability, and mental health issues are all forecasted. Communities that will be especially vulnerable will be: rural and low-income communities, BIPOC communities, private well users, agricultural and outdoor recreation workers, firefighters and other first responders, and children and pregnant women.

Mid-Oregon Coastal Concerns

Mote *et al.* (2019) report that coastal Oregon can expect more severe winter storms, increased ocean temperature is likely to produce harmful algal blooms affecting commercial, recreational, and tribal fisheries. Meanwhile, ocean acidification will compromise fisheries, especially shellfish, and sea level rise, plausibly reaching over 8 feet by 2100, will increase the risk of flooding, coastal erosion and threaten estuaries and coastal marshes (Mote *et al.* 2019).

Because much of Oregon's 5th Senate District lies along the state's famed coastline, this district has special concerns to address as climate change progresses. One result of climate change is a rise in sea level. If sea levels continue to rise throughout this and the next century, then Oregon's 5th Senate District could face development and infrastructure problems in its coastal cities. The 5th Senate District's economy relies heavily on tourism generated by these popular vacation spots near the coast, and much of this revenue could be lost if nothing is done to buffer coastal cities from the negative impacts of climate change. Predictions for coastal Oregon range from 6 inches over the next century to a sea level rise of 8 feet (Mote *et al.* 2019). Even a sea level rise of 6 inches could be devastating to coastal Oregon's infrastructure, economic development, and the booming tourism industry.

The port of Coos Bay, previously the largest shipper of timber in the world, still plays an important part in the region's economy. However, coastal regions are particularly threatened by climate change. Some sources predict that sea levels on the Oregon coast could rise anywhere from 5" to 6 feet. A rise of even 5" could be devastating to coastal infrastructure and development, especially coastal ports. Such damage would also undoubtedly affect the tourism industry along Oregon's coast.

Coastal regions face other challenges, as well, as climate change continues to progress. Studies have shown that ocean acidification, a direct consequence of rising carbon levels from carbon emissions, can have devastating consequences to the development of juvenile fish and the survival of adult fish. Ocean acidification is undoubtedly going to negatively impact coastal Oregon's fishing industry, as fish become more and more scarce.

Because of scientific models, we can predict when and how climate change is going to affect Oregon's 5th Senate District. While we may not be able to reverse climate change, we can take steps to ensure that Oregon's 5th Senate District's economy continues to thrive, even as the climate changes. It is vital, however, that we act now in order to mitigate problems before they arise.

A Timeline for Action:

Based on the projected consequences of the warming global climate, international agreements (e.g., UNFCCC 2015) some years ago established 2°C (preferably 1.5°C) above pre-industrial conditions as the limit beyond which we should not allow the global temperature to climb. This limit was echoed by the World Bank (2014). Meanwhile, the Intergovernmental Panel on Climate Change (IPCC 2018a) indicated that the 2°C limit pushes us too close to many global tipping points beyond which recovery becomes a reducing possibility. Thus, they recommend that we absolutely should target 1.5°C if we wish a reasonable chance of retaining a livable planet. Unfortunately, underlining the urgency, emissions to date may have already committed us to the 1.5°C increase (Mauritsen and Pincus 2017).

Global greenhouse gas emissions during 2017 totaled 53.7 Gigatonnes (GT) of carbon dioxide equivalent (IPCC 2018b) which includes between 32.5 (IEA 2019) and 36.5 GT of carbon dioxide (WRI 2018). This implies that between 30% and 40% of the global warming emissions are due to gases other than carbon dioxide. The trends and consequences discussed here are based on readily available data. This underlines the urgency for immediate action across the globe to curtail greenhouse gas emissions if we wish to avoid an increase over 2°C. Considerable variability exists among estimates of the emissions budget remaining if we are to restrict warming to the 1.5°C increase targets (Levin 2018, Carbon Brief 2018). Indeed, the latter source identifies a large range in estimates for a 66% chance of keeping warming to below 1.5°C of between 28 GT and 779 GT. Meanwhile, the IPCC (2018a) indicated that the rate of carbon dioxide emissions alone is currently 42 ± 3 Gigatonnes annually suggesting that, for a 50% chance at a rise below 1.5°C, the remaining budget for emissions is 580 GT CO₂, while for a 66% chance, the remaining emissions budget is 420 GT CO₂. Considering the current accelerating rate of emissions, the IPCC (2018a) concluded that by 2030 we must impose a reduction in emissions of 45% below the 2010 level and by 2050 we must reach net zero emissions. Considering the increasing impact of greenhouse gases other than carbon dioxide, that seems both conservative and reasonable. Underlining the urgency and imperative of limiting warming to 1.5°C, long ago the World Bank (2014) acknowledged there is: "no certainty that adaptation to a 4°C world is possible."

Representing the People at the 24th United Nations Framework Convention on Climate Change Conference of the Parties in Poland (COP24), British naturalist and broadcaster Sir David Attenborough argued that in climate change “we are facing a man-made disaster of global scale, our greatest threat in thousands of years...” and “If we don't take action, the collapse of our civilizations and the extinction of much of the natural world is on the horizon.” (Domonoske 2018). The choice is ours!

From the trends and consequences discussed here, all based on readily available data, there should be little doubt that substantial urgency must be attached to addressing this issue.

Solutions:

In addition to individual action wherein we evaluate our actions and adjust our behavior to reduce activities that result in greenhouse gas emissions or increase those that result in atmospheric greenhouse gas sequestration (capture and storage), we can promote local, state and federal actions that do the same on a larger scale. Local communities can develop Climate Action Plans that promote emissions reductions and capture/sequestration activities. Meanwhile, at the state and federal level, similar such programs can be instituted. The predominant proposals to achieve this involve either:

- a) imposing a jurisdictional cap on emissions which declines over time to establish a trajectory of emissions reductions that meet long term reductions goals. This approach involves the issuance of allowances to emit that reduce over time. Allowances may be sold/auctioned, or allocated free, or involve some combination.
- b) imposing a fee or tax on emissions that rises over time to achieve reductions that are consistent with a desired trajectory and long-term goals.

The cap approach is direct since it involves assessing emissions from target polluters and requiring that reductions occur while the tax/fee approach is indirect since it is based on the assumption that a rising tax will result in reduced emissions.

Both approaches usually involve the generation of funds either via sold/auctioned allowance in the case of the cap, or a fee in the case of the tax/fee approach. The second question associated with either approach involves a decision as to what will be done with the funds raised. One approach is to return these to residents or taxpayers (the individuals who ultimately pay the cost of the pollution reduction); hence the concept of a Dividend. Alternatively, the funds raised can be used to offset allow reductions in other taxes, whether individual or corporate. Finally, these funds may be used for investments that (a) promote activities that themselves lead to reductions in atmospheric greenhouse gas concentrations, either by reducing emissions or promoting sequestration and/or (b) serve the goals of

promoting environmental / social justice by assisting communities historically disadvantaged by pollution or likely to suffer disproportionately from the transition to a clean energy economy.

Since the state legislature has declined to implement a comprehensive policy, if Oregon is to contribute its share to addressing the climate crisis and wishes to appear credible when seeking action elsewhere, it will be necessary to take smaller targeted steps that reduce emissions in designated sectors or activities and/or promote the sequestration of carbon from our atmosphere in our natural and working lands.

The Potential of Offshore Wind:

With more than 7,500 miles of coastline, including Hawaii and Alaska, the Pacific Coast of the United States holds enormous potential for the extraction of marine renewable energy. The region has capable ports along the length of its coastline as well as abundant capacity within the electrical grid. The proximity of manufacturing operations to the coastline and well-developed transportation systems makes it optimal for early-stage technology developers looking to test and develop in the same location (Poet 2023). Two locations in Senate District 1 have been identified as high potential areas for offshore wind development. The Coos Bay and Brookings areas have both been targeted for this kind of development and there are currently groups working to establish this source of energy (Parks 2022). It will take a concerted effort by advocates and decision makers to bring this potential source of sustainable energy and jobs to fruition.

Contact Your Legislators:

Contact Senator Dick Anderson

Capitol Phone: 503-986-1705

Capitol Address: 900 Court St NE, S-303, Salem, OR, 97301

Email: Sen.DickAnderson@oregonlegislature.gov

Website: <http://www.oregonlegislature.gov/anderson>

House District 9: Boomer Wright

Capitol Phone: 503-986-1409

Capitol Address: 900 Court St NE, H-372, Salem, OR 97301

Email: Rep.BoomerWright@oregonlegislature.gov

Website: <http://www.oregonlegislature.gov/wright>

House District 10: David Gomberg

Capitol Phone: 503-986-1410

Capitol Address: 900 Court St NE, H-471, Salem, OR 97301

Email: Rep.DavidGomberg@state.or.us

Website: <http://www.oregonlegislature.gov/gomberg>

Literature:

Alexander M, Scott J, Friedland K, Mills K, Nye J, Pershing A, Thomas A, 2018. Projected sea surface temperatures over the 21st century: Changes in the mean, variability and extremes for large marine ecosystem regions of Northern Oceans. *Elem Sci Anth*, 6(1), p.9. DOI:

<http://doi.org/10.1525/elementa.191>

Bednaršek N, Feely R, Beck M, Alin S, Siedlecki S, Calosi P, Norton E, Caenger C, Strus J, Greeley D, Nezlin N, Roethler M, Spicer J. 2020 Exoskeleton dissolution with mechanoreceptor damage in larval Dungeness crab related to severity of present-day ocean acidification vertical gradients. *Science of the Total Environment*. 716, 10 136610

<https://doi.org/10.1016/j.scitotenv.2020.136610>

Ben Achur S. 2022 Ocean acidification raises economic concerns for shellfish hatcheries. Market Place Minnesota Public Radio July 5 2022. <https://www.marketplace.org/2022/07/05/ocean-acidification-raises-economic-concerns-shellfish-hatcheries/>

Brown L 2006 *Plan B 2.0: Rescuing a Planet Under Stress and a Civilization in Trouble*. W.W. Norton, & Co. N.Y. London 365 pp.

Carbon Brief 2018 Analysis: How much ‘carbon budget’ is left to limit global warming to 1.5°C?

<https://www.carbonbrief.org/analysis-how-much-carbon-budget-is-left-to-limit-global-warming-to-1-5c>

Crookston N, Radtke P. 2023 Plant Species and Climate Profile Predictions, Climate Estimates, Climate Change and Plant Climate Relationships.

<https://charcoal2.cnre.vt.edu//climate/species/index.php> (A discussion of the site can be found at: <https://www.fs.usda.gov/ccrc/tool/plant-species-and-climate-profile-predictions> though the site has been relocated from the USDA so the links on that page are dead ends).

CTT 2021 Congressional Temperature Trends: Thermometer Records from Weather Stations

<http://temperaturetrends.org/district.php?district=4&state=OR>

Dalton M, Mote P, Snover A [Eds.]. 2013. *Climate Change in the Northwest: Implications for Our Landscapes, Waters, and Communities*. Washington, DC: Island Press.

<http://cses.washington.edu/db/pdf/daltonetal678.pdf>

Dalton M, Dello K, Hawkins L, Mote P, and Rupp D (2017) The Third Oregon Climate Assessment Report, Oregon Climate Change Research Institute, College of Earth, Ocean and Atmospheric Sciences, Oregon State University, Corvallis, OR.

http://www.occri.net/media/1055/ocar3_final_all_01-30-2017_compressed.pdf

Domonoske C, 2018 David Attenborough Warns Of 'Collapse Of Civilizations' At U.N. Climate Meeting. npr news. <https://www.npr.org/2018/12/03/672893695/david-attenborough-warns-of-collapse-of-civilizations-at-u-n-climate-meeting>

Easterling, D, Kunkel K, Arnold J, Knutson T, LeGrande A, Leung L, Vose R, Waliser D, Wehner M, 2017: Precipitation change in the United States. In: Climate Science Special Report:Fourth National Climate Assessment, Volume I [Wuebbles, D, Fahey D, Hibbard K, Dokken D, Stewart B, Maycock T. (eds.)]. U.S. Global Change Research Program, Washington, DC, USA, pp. 207-230, doi: 10.7930/J0H993CC.

https://science2017.globalchange.gov/downloads/CSSR_Ch7_Precipitation.pdf

Fleishman, E., editor. 2023. Sixth Oregon Climate Assessment. Oregon Climate Change Research Institute, Oregon State University, Corvallis, Oregon.

<https://blogs.oregonstate.edu/occri/oregon-climate-assessments>

Gao P, Terando A, Kupfer J, Varner J, Stambaugh M, Lei T, Hiers J. 2021 Robust projections of future fire probability for the conterminous United States. *Science of the Total Environment* 789: 147872. <https://www.sciencedirect.com/science/article/pii/S0048969721029430>

Halofsky J, Peterson D, Metlen K, Myer G, Sample V. 2016 Developing and Implementing Climate Change Adaptation Options in Forest Ecosystems: A Case Study in Southwestern Oregon, USA forests 7: (268) 1 - 18.

<file:///F:/Alan/Documents/KACONJOUR/Climate%20%20Change%20Literature/Forests,%20Fire%20&%20BLM%20RMP/Halofsky,%20Peterson,%20%20Metlen,%20Myer,%20Sample%202016%20climate%20change%20and%20adaptation%20in%20the%20Rogue%20Basin%20Oregon.pdf>

Hausfather Z. 2018 Explainer: How 'Shared Socioeconomic Pathways' explore future climate change. Climate Modeling. Carbon Brief. <https://www.carbonbrief.org/explainer-how-shared-socioeconomic-pathways-explore-future-climate-change/>

Howat I, Tulascyk S. 2005 Trends in spring snowpack over a half-century of climate warming in California, USA *Annals of Glaciology* 40: 151-156.

<https://www.cambridge.org/core/journals/annals-of-glaciology/article/trends-in-spring-snowpack-over-a-halfcentury-of-climate-warming-in-california-usa/E7153C040C54C29DFC9EDB92678C02FD>

Hudiburg T, Law B, Turner D, Campbell J, Donata D, Duane M. 2009 Carbon dynamics of Oregon and Northern California forests and potential land-based carbon storage. Ecological Applications. <https://doi.org/10.1890/07-2006.1>

IPCC 2013. Climate Change 2013: The Physical Science Basis; Summary for Policymakers WGI IPCC Switzerland. <https://www.ipcc.ch/report/ar5/wg1/>

IPCC 2018a. Global Warming of 1.5°C, (An IPCC Special Report) Intergovernmental Panel on Climate Change Summary for Policymakers - https://report.ipcc.ch/sr15/pdf/sr15_spm_final.pdf
Full Report - <https://www.ipcc.ch/sr15/>

IPCC 2018b. Emissions Gap Report United Nations Environment Programme
<https://www.ipcc.ch/site/assets/uploads/2018/12/UNEP-1.pdf>

IUCN undated. Overview and description. IUCN CEM Mediterranean Type Ecosystems Specialist Group. International Union for the Conservation of Nature. [https://www.iucn.org/our-union/commissions/group/iucn-cem-mediterranean-type-ecosystems-specialist-group#:~:text=Mediterranean%2Dtype%20ecosystems%20\(MTEs\),and%20Southwestern%20and%20South%20Australia](https://www.iucn.org/our-union/commissions/group/iucn-cem-mediterranean-type-ecosystems-specialist-group#:~:text=Mediterranean%2Dtype%20ecosystems%20(MTEs),and%20Southwestern%20and%20South%20Australia).

Jones G 2015 Climate, Grapes and Wine: Terroir and the Importance of Climate to Winegrape Production. GuildSomm
https://www.guildsomm.com/public_content/features/articles/b/gregory_jones/posts/climate-grapes-and-wine

Jones N 2017 How the World Passed a Carbon Threshold and Why It Matters. Yale Environment 350 <https://e360.yale.edu/features/how-the-world-passed-a-carbon-threshold-400ppm-and-why-it-matters>

Karl T, Melillo J, Peterson T 2009 Global Climate Change Impacts in the United States, Cambridge University Press <https://downloads.globalchange.gov/usimpacts/pdfs/climate-impacts-report.pdf>

Kenward A, Sanford T, Bronzan J 2016 Western Wildfires: A Fiery Future. Climate Central: <file:///H:/Alan/Documents/KACONJOUR/Climate%20Change%20literature/Forests,%20Fire%20&%20BLM%20RMP/Kenward%20et%20al%20westernwildfires2016vfinal.pdf>

Law B, Hudiburg T, Berner L, Harmon M 2018 Land use strategies to mitigate climate change in carbon dense temperate forests. Proceedings of the National Academy of Sciences. 115 (14) 3663-3668. <https://www.pnas.org/doi/10.1073/pnas.1720064115>

Levin K 2018 According to New IPCC Report, the World Is on Track to Exceed its “Carbon Budget” in 12 Years, World resources Institute. <https://www.wri.org/blog/2018/10/according-new-ipcc-report-world-track-exceed-its-carbon-budget-12-years>

Lieberman B 2012 ‘CSZ’ — A Key Factor for Pacific Northwest Sea-Level Rise. Yale Climate Connections. <https://www.yaleclimateconnections.org/2012/07/cascadia-subduction-zone-a-key-factor-for-pacific-nw-sea-level-rise/>

Maycock, and B.C. Stewart (eds.). U.S. Global Change Research Program, Washington, DC, USA, 1515 pp. doi: 10.7930/NCA4.2018 <https://nca2018.globalchange.gov/>

Melillo, J, Richmond T, Yohe G, Eds., 2014: Climate Change Impacts in the United States: The Third National Climate Assessment. U.S. Global Change Research Program, 841 pp. doi:10.7930/J0Z31WJ2. http://s3.amazonaws.com/nca2014/low/NCA3_Climate_Change_Impacts_in_the_United%20States_LowRes.pdf?download=1

Mote, P.W., J. Abatzoglou, K.D. Dello, K. Hegewisch, and D.E. Rupp, 2019: Fourth Oregon Climate Assessment Report. Oregon Climate Change Research Institute. occri.net/ocar4. <http://www.occri.net/media/1095/ocar4full.pdf>

NASA GISS 2023 Annual Mean Temperature Change Over Land and Over Ocean. GISS Surface Temperature Analysis (v4). National Aeronautics and Space Administration Goddard Institute for Space Studies. https://data.giss.nasa.gov/gistemp/graphs_v4/

NOAA 2023 Climate At A Glance National Time Series. National Oceanographic and Atmospheric Administration. https://www.ncei.noaa.gov/access/monitoring/climate-at-a-glance/national/time-series/110/tavg/12/12/1895-2022?base_prd=true&begbaseyear=1901&endbaseyear=2000&trend=true&trend_base=100&begtrendyear=1895&endtrendyear=2023&filter=true&filterType=binomial

Oregon Health Authority 2014, Oregon Climate and Health Profile Report, Climate and Health Program, Public Health Division, Oregon Health Authority: Summary provided by Emily York MPH, Climate & Health Program Coordinator, Oregon Public Health Division, Oregon Health Authority. <http://public.health.oregon.gov/HealthyEnvironments/climatechange/Documents/oregon-climate-and-health-profile-report.pdf>

Parks B 2022. Oregon Public Broadcasting. <https://www.opb.org/article/2022/04/27/offshore-wind-oregon-coast-coos-bay-brookings/>

Poet 2023 Pacific Coast Advantage. Pacific Ocean Energy Trust: <https://pacificoceanenergy.org/information/pac-coast-advantage/>.

Rehfeldt G and Crookston N. 2023 Plant Species and Climate Profile Predictions, <http://charcoal2.cnre.vt.edu/climate> (regrettably as of 02/08/2023 this site is under repair from a hack). A discussion of the site can be found at: <https://www.fs.usda.gov/ccrc/tool/plant-species-and-climate-profile-predictions> though the site has been relocated from the USDA so the links on that page are dead ends. UNFCCC 2015. United Nations Framework Convention on Climate Change Adoption of the Paris Agreement 32pp <https://unfccc.int/resource/docs/2015/cop21/eng/l09r01.pdf>

Safford H, Butz R, Bohlman G, Coppoletta M, Estes B, Gross S, Merriam K, Meyer M, Molinari N, Wuenschel A. 2021 Fire Ecology of the North American Mediterranean-Climate Zone. Chapter 9 https://www.fs.usda.gov/psw/publications/srs/srs_2021_safford001.pdf [In] Greenberg C, Collins B (eds.), Fire Ecology and Management: Past, Present, and Future of US Forested Ecosystems, Managing Forest Ecosystems 39, https://doi.org/10.1007/978-3-030-73267-7_9.

SOS 2022 Climate Model - Surface Temperature Change: SSP1 (Sustainability) - 2015 – 2100. Science on a Sphere. National Oceanographic and Atmospheric Administration <https://sos.noaa.gov/catalog/datasets/climate-model-surface-temperature-change-ssp1-sustainability-2015-2100/>

UNFCCC 2015 The Paris Agreement Unites Nations Climate Change. <https://unfccc.int/process-and-meetings/the-paris-agreement/the-paris-agreement>

USGCRP, 2017: Climate Science Special Report: Fourth National Climate Assessment, Volume I [Wuebbles D, Fahey D, Hibbard K, Dokken D, Stewart B, and Maycock T (eds.)]. U.S. Global Change Research Program, Washington, DC, USA, 470 pp <https://science2017.globalchange.gov/downloads/>

USGS 2019 National Climate Change Viewer, U.S. Geological Survey Land Change Science Program. <https://www2.usgs.gov/landresources/lcs/nccv/viewer.asp>

Westerling A, Hidalgo H, Cayan D, Swetnam D. 2006 Warming and Earlier Spring Increase Western U.S. Forest Wildfire Activity. *Science* 313 no. 5789 pp. 940-943.

<https://www.science.org/doi/10.1126/science.1128834>

Whittaker RH 1975 *Communities and Ecosystems* Macmillan Publishing CO., New York

World bank 2014, *Turn Down the Heat: Confronting the New Climate Normal*. The World Bank, Washington DC. 275pp <https://openknowledge.worldbank.org/entities/publication/98508814-21c5-53e6-b36c-912a4ecf9da7>

WRI 2018 New Global CO2 Emissions Numbers Are In. They're Not Good. World Resources Institute. <https://www.wri.org/blog/2018/12/new-global-co2-emissions-numbers-are-they-re-not-good>